

Table (1): Vital signs among the participants

Variable	Mean ± SD (Range)
Systolic blood pressure, mmhg	137 ± 17 (120 – 200)
Diastolic blood pressure, mmhg	87 ± 6.1 (80 – 110)
Pulse, bpm	95 ± 12 (80 – 140)
Temperature, degrees	37.5 ± 0.2 (36.5 – 38)
BMI, kg/m²	32.1 ± 5.7 (20 – 40)

Table (2): Fetal characters among the participants

Variable	No (%)
Fetal presentation	Cephalic 58 (58)
	Breech 32 (32)
	Other 10 (10)
IUGR, n (%)	10 (10)
Congenital anomalies, n (%)	8 (8)
Estimated fetal weight (mean±SD (Range))	3070 ± 607 (1571 – 4000)
Gestational age (mean±SD (Range))	37.6 ± 1.2 (34 – 39)

Table (3): Feto-maternal outcome among the participants

Variable	No (%)	
Preterm delivery, n (%)	14 (14)	
Mode of delivery	Vaginal	48 (48)
	CS	52 (52)
Weight at delivery, mean± SD	3227.4± 639.5	
Gender	Male	55 (55)
	Female	45 (45)
1-min APGAR<7	23 (23)	
5-min APGAR<7	14 (14)	
NICU, n (%)	14 (14)	
NICU duration	<24 hours	6 (6)
	24-48 hours	5 (5)
	> 48 hours	3 (3)
RDS, n (%)	7 (7)	
Neonatal mortality, n (%)	5 (5)	

RDS; Respiratory distress syndrome

Table (4): Comparison between the three degrees of polyhydramnios regarding feto-maternal outcome

Variable	Mild (n = 46)	Moderate (n = 28)	Severe (n = 26)	P-value
IUGR	3 (6.5)	6 (21.4)	1 (3.8)	0.056
Preterm delivery. N (%)	9 (19.6)	1 (3.6)	4 (15.4)	0.153
Mode of delivery	Vaginal	15 (53.6)	14 (53.8)	0.785
	CS	25 (54.3)	13 (46.4)	
Weight at delivery, mean± SD	3202 ± 640	3307 ± 573	3185 ± 717	0.739
1-min APGAR<7	12 (26.1)	8 (28.6)	3 (11.5)	0.264
5-min APGAR<7	8 (17.4)	6 (21.4)	3 (11.5)	0.624
NICU, n (%)	10 (21.7)	5 (17.9)	5 (19.2)	0.915
NICU duration	<24 hours	5 (10.9)	0 (0)	0.634
	24-48 hours	1 (2.2)	2 (7.1)	
	> 48 hours	0 (0)	1 (3.6)	
RDS, n (%)	5 (10.9)	1 (3.6)	1 (3.8)	0.375
Neonatal mortality, n (%)	1 (2.2)	2 (7.1)	2 (7.6)	0.553