

**Table (1): Distribution of demographic characteristics among the studied women**

Parameters	Studied women (n=46)		
	N	%	
<b>Residence</b>	<b>Al-Minya</b>	36	78.2%
	<b>Assuit</b>	1	2.2%
	<b>Bani-Suweif</b>	7	15.2%
	<b>Sohag</b>	1	2.2%
	<b>Cairo</b>	1	2.2%
<b>Age (years)</b>	<b>Mean± SD</b>	30.43± 6.12	
	<b>Median</b>	30.0	
	<b>Range</b>	18.0- 43.0	
	<b>18-28</b>	17	37%
	<b>29-38</b>	24	52.2%
	<b>&gt; 39</b>	5	10.8%
<b>BMI (Kg/m<sup>2</sup>)</b>	<b>Mean± SD</b>	23.04± 2.18	
	<b>Median</b>	23.0	
	<b>Range</b>	19.4 – 26.5	
<b>Educational level</b>	<b>Illiterate</b>	4	8.7%
	<b>Read&amp;Write</b>	6	13%
	<b>Primary Education</b>	8	17.4%
	<b>Preparatory Education</b>	9	19.6%
	<b>Secondary Education</b>	13	28.3%
	<b>High Education</b>	6	13%

*SD= standard deviation,*

**Table (2): Distribution of diagnosis period among the studied women**

Parameters	Studied women (n=46)	
	N	%
Diagnosis period	1 <sup>st</sup> trimester	3 6.5%
	2 <sup>nd</sup> trimester	3 6.5%
	3 <sup>rd</sup> trimester	35 76.1%
	Labour	5 10.9%

**Table (3): Distribution of clinical presentation among the studied women**

Parameters	Studied women (n=46)	
	N	%
Asymptomatic	1	2.2%
Fever	40	87.0%
Dyspnea	26	56.5%
Tachypnea	5	10.9%
Cough	41	89.1%
Hypoxia	4	8.7%
PROM	4	8.7%
Sore throat	6	13.0%
Labor pain	10	21.7%
Tender Scar	2	4.3%
Vomiting	2	4.3%
Hemoptysis	1	2.2%
Tachycardia	1	2.2%
Diarrhea	1	2.2%
Hypertension	3	6.5%
Palpitation	1	2.2%
Abdominal Pain	2	4.3%
Vaginal Bleeding	5	10.9%
Pallor	1	2.2%
Pelvic Pain	2	4.3%
Combined	45	97.8%

**Table (4): Distribution of clinical data among the studied women**

Parameters	Studied women (n=46)	
	N	%
Induction of labor	No	45 97.8%
	Yes	1 2.2%
Hospital stay (days)	Mean± SD	10.04± 6.04
	Median	8.0
	Range	3.0- 30.0
42-day readmission	No	44 95.7%
	Yes	2 4.3%
Mode of delivery	NVD	10 21.7%
	C.S	35 76.1%

**Table (5): Distribution of maternal complications among the studied women**

Parameters	Studied women (n=46)	
	N	%
Non-complicated	29	63%
Maternal complications	ICU admission	17 37.0%
	Mechanical ventilation	7 15.2%
	Maternal death	7 15.9%

**Table (6): Maternal death among the studied women**

Parameters	Studied women (n=46)			
	N	%		
<b>NO of dead women</b>	<b>Total women = 46</b>	7	15.9%	
<b>Residence</b>	<b>El-Minia</b>	7	100%	
<b>Age (years)</b>	<b>Mean± SD</b>	32.3±5.68		
	<b>Median</b>	31		
	<b>Range</b>	27.0 – 39.0		
	<b>25-30</b>	2	28.6%	
	<b>30 - 35</b>	3	42.8 %	
	<b>35 -40</b>	2	28.6%	
<b>BMI (Kg/m<sup>2</sup>)</b>	<b>Mean± SD</b>	23.04± 4.8		
	<b>Median</b>	23.2		
	<b>Range</b>	21.1– 24.8		
<b>Educational level</b>	<b>Read &amp; Write</b>	1	14.3%	
	<b>Primary Education</b>	1	14.3%	
	<b>Preparatory Edu.</b>	2	28.6%	
	<b>Secondary Edu.</b>	2	28.6%	
	<b>High Education</b>	1	14.2%	
	<b>Clinical Presentation</b>	<b>Cough</b>	7	100%
<b>Fever</b>		6	85.7%	
<b>Dyspnea</b>		6	85.7%	
<b>Tachypnea</b>		2	28.6%	
<b>Hypertension</b>		2	28.6%	
<b>Vaginal bleeding</b>		2	28.6%	
<b>Abdominal pain</b>		1	14.3%	
<b>Hypoxia</b>		1	14.3%	
<b>Combined</b>		7	100%	
<b>Obstetric History</b>	<b>Primi Gravida</b>	2	28.6%	
	<b>Multi Gravida</b>	5	71.4%	
	<b>Gestational Age (weeks)</b>	<b>Mean± SD</b>	34± 5	
		<b>Median</b>	33	
		<b>Range</b>	29 -37	
<b>Associated Co- Morbidity</b>	<b>Non Associated</b>	1	14.3%	
	<b>Gestational Hypertension</b>	2	28.6%	
	<b>Gestational Diabetes Mellitus</b>	2	28.6%	
	<b>Placenta Previa</b>	2	28.6%	
	<b>Rupture uterus</b>	1	14.3%	
	<b>Cardiac lesion</b>	1	14.3%	
	<b>Chronic Hypertension</b>	1	14.3%	
	<b>Eclampsia</b>	1	14.3%	

**Table (7): Distribution of neonatal morbidity & mortality among the studied women**

<b>Parameters</b>	<b>Studied cases (n=46)</b>	
	<b>N</b>	<b>%</b>
<b>Neonatal complications</b>	<b>Non complicated</b>	24 52.2%
	<b>Preterm</b>	6 13.0%
	<b>Neonatal jaundice</b>	1 2.2%
	<b>RDS</b>	4 8.7%
	<b>IUFD</b>	5 10.9%
	<b>Missed abortion</b>	3 6.5%
	<b>Died after labour</b>	1 2.2%